

THE COMPLEX GRADIENT INEQUALITY WITH PARAMETER

MACIEJ P. DENKOWSKI

ABSTRACT. We prove that given a holomorphic family of holomorphic functions with isolated singularities at zero and constant Milnor number, it is possible to obtain the gradient inequality with a uniform exponent.

1. INTRODUCTION

This note was inspired by a recent article of A. Płoski [P2] concerning the semi-continuity of the Łojasiewicz exponent in a family of multiplicity-constant deformation of a finite holomorphic germ (actually, our main tool is a method of Płoski used already in [P1]). Of course, it would probably be possible to derive the result presented herein from this article, but it seems interesting to give a direct, elementary and self-consistent proof, since it is a starting point for tackling a much more general problem.

We are interested in the Łojasiewicz Gradient Inequality: given an (complex or real) analytic (or subanalytic and of class $\mathcal{C}^1$) function f of m variables and such that $f(0) = 0$ we are able to find an exponent $\theta \in (0, 1)$ such that in a neighbourhood of zero

$$|f(x)|^\theta \leq \text{const.} \|\nabla f(x)\|$$

where $\nabla f(x)$ denotes the (complex or real) gradient of f . The classical version (for a real analytic function) was obtained by Łojasiewicz from his famous Łojasiewicz inequality that he used to solve L. Schwartz's Division Problem. One of the simplest and most important applications of this inequality is the study of the analytic gradient dynamics, see [L] (the fact that $\theta \in (0, 1)$ is of utmost importance; to make the paper self-consistent, we give a simple proof of this in our special case in Section 2).

Our general aim is to obtain a parameter version of this inequality, i.e. to prove that given a well-parametrized family of functions we can find (locally) a uniform exponent for this inequality to be satisfied by each of these functions. Probably the simplest case is presented in this paper and it concerns a μ -constant holomorphic unfolding.

Date: March 4th 2014.

1991 Mathematics Subject Classification. 32B20.

Key words and phrases. Gradient inequality, holomorphic functions.

A general subanalytic parameter version of the gradient inequality will be given in a forthcoming paper.

2. ON THE GRADIENT EXPONENT

For convenience sake, we will give — using only complex analytic geometry tools in the spirit of Płoski — yet another and elementary proof of the gradient inequality for a holomorphic germ $f: (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}, 0)$ satisfying the condition $\dim_0 \nabla f^{-1}(0) = 0$, where

$$\nabla f = \left(\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_m} \right) : \mathbb{C}_z^m \rightarrow \mathbb{C}_w^m$$

is the complex gradient. What will be of greatest interest for our purposes is the bound on the exponent we obtain.

We may assume $m > 1$, since the case $m = 1$ is a simple exercise involving power series. Hereafter we will insist on details that for simplicity we omit in the next section.

Our approach is based on [P1]. Take a connected neighbourhood U of $0 \in \mathbb{C}^m$ for which $\nabla f^{-1}(0) \cap U = \nabla f^{-1}(0) \cap \overline{U} = \{0\}$. Hence ∇f is a proper map from U onto a connected neighbourhood V of zero. This means that the projection $\pi(z, w) = w$ is proper on $\Gamma_{\nabla f} \cap U \times V$ and sends this part of the graph onto V . It is thence a branched covering (see [Ch]) with covering number μ . Its singular locus¹ $\sigma \subsetneq V$ is analytic and μ is in fact the *Milnor number* of f at zero. For $w \in V \setminus \sigma$ we have $\nabla f^{-1}(w) = \{z^{(1)}, \dots, z^{(\mu)}\}$ consisting of exactly μ points and so we can define for $t \in \mathbb{C}$

$$\begin{aligned} P(w, t) &:= \prod_{z \in \nabla f^{-1}(w)} (t - f(z)) = \\ &= t^\mu + a_1(w)t^{\mu-1} + \dots + a_\mu(w), \\ \text{where } a_j(w) &= (-1)^j \sum_{1 \leq \iota_1 < \dots < \iota_j \leq \mu} f(z^{(\iota_1)}) \cdot \dots \cdot f(z^{(\iota_j)}) \end{aligned}$$

are holomorphic functions on $V \setminus \sigma$ admitting continuous extensions onto V . Therefore, by the Riemann Extension Theorem, $P \in \mathcal{O}(V)[t]$. This polynomial is called *the characteristic polynomial of f* with respect to ∇f . Clearly, $P(\nabla f, f) \equiv 0$ and $a_j(0) = 0$.

Remark 2.1. Note that in the same way we can define the characteristic polynomial of f with respect to any holomorphic map germ $g: (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ satisfying $g^{-1}(0) = \{0\}$.

The Viète formulæ yield $|f(z)| \leq 2 \max_j |a_j(\nabla f(z))|^{1/j}$. On the other hand, since in a neighbourhood of zero there is $|a_j(w)| \leq c_j \|w\|^{\text{ord}_0 a_j}$

¹i.e. the set of points $w \in V$ for which $\#\nabla f^{-1}(w) < \mu$.

for some $c_j > 0$ ($\text{ord}_0 h$ denotes the order of vanishing at zero² of the holomorphic function h), this leads to

$$|f(z)| \leq \text{const.} \|\nabla f(z)\|^{\min_{j=1}^{\mu} \left(\frac{\text{ord}_0 a_j}{j} \right)}$$

in a neighbourhood of zero. We have obtained the gradient inequality with the exponent

$$\theta = \max_{j=1}^{\mu} \frac{j}{\text{ord}_0 a_j}$$

and we shall prove in addition that $\theta < 1$ (i.e. $\text{ord}_0 a_j > j$ for all j). This is still rather delicate (see the crucial article [T] of B. Teissier, Corollaire 2 p. 270 and compare with [PW] Theorem 2).

Geometrically, the weaker condition:

$$\text{ord}_0 a_j \geq j \text{ for all } j,$$

is equivalent to saying that the tangent cone of $P^{-1}(0)$ at zero, described by the zeroes of the initial form in P in the expansion of P into homogeneous forms at zero meets $\{0\}^m \times \mathbb{C}$ only at zero (since this means precisely that $\text{in}P(0, t) = t^\mu$; actually, we will show that $\text{in}P(w, t) = t^\mu$). Recall the Peano tangent cone:

$$C_0(P^{-1}(0)) = \{v \in \mathbb{C}^{m+1} \mid \exists P^{-1}(0) \ni x_\nu \rightarrow 0, \exists \lambda_\nu > 0: \lambda_\nu x_\nu \rightarrow v\}.$$

The weak version follows from [P1], as $P(\nabla f, f) = 0$ is an equation of integral dependence of f over the ideal $\left\langle \frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_m} \right\rangle$ which implies (actually, is equivalent to) the inequality

$$(\star) \quad |f(z)| \leq \text{const.} \|\nabla f(z)\|$$

in a neighbourhood of zero³. Now, since $P^{-1}(0)$ and $(\nabla f, f)(U)$ define the same set germ at zero, we see that for any $v = (0, t) \in C_0(P^{-1}(0))$ we can find sequences $U \ni z_\nu \rightarrow 0$ and $\lambda_\nu \rightarrow 0^+$ such that $\lambda_\nu \nabla f(z_\nu) \rightarrow 0$ and $\lambda_\nu f(z_\nu) \rightarrow t$. Therefore, $(\star)$ implies $t = 0$. We need only to prove that $\mathbb{C}^m \times \{0\} \supset C_0(P^{-1}(0))$ (for we know that $C_0(P^{-1}(0))$ is an algebraic cone of dimension $\dim_0 P^{-1}(0) = m$). We have just seen that $C_0(P^{-1}(0)) \cap \{0\}^m \times \mathbb{C} = \{0\}^{m+1}$. Shrinking the neighbourhoods we may assume that $P^{-1}(0) = (\nabla f, f)(U)$ and this has pure dimension m . Then we can use the elementary Proposition from [Ch] p. 86: for any $(w, t) \in C_0(P^{-1}(0)) \setminus \{(0, 0)\}$ (in particular for $w \neq 0$) we have

$$(w, t) \in C_0(P^{-1}(0) \cap (\mathbb{C}w \times \mathbb{C})) =: C.$$

² $\text{ord}_0 h$ is the degree of the initial form in the expansion of $h \neq 0$ into homogeneous forms near zero. There is $\text{ord}_0 h = \max\{\eta > 0 \mid |h(z)| \leq \text{const.} \|z\|^\eta \text{ in a neighbourhood of } 0\}$, see [Ch].

³From the expressions for a_j and the Viète formulæ it is easy to see that $P(\nabla f, f) \equiv 0$ implies $|f(z)| \leq \text{const.} \|\nabla f(z)\|$. For the other way round it suffices to observe that a_j belongs to $\mathfrak{m}^j = \langle z^\alpha: |\alpha| = j \rangle$, where $\mathfrak{m}$ is the maximal ideal of the ring $\mathcal{O}_m$.

But, $P^{-1}(0) \cap (\mathbb{C}w \times \mathbb{C}) = \{(\nabla f(z), f(z)) \mid z \in \nabla f^{-1}(\mathbb{C}w)\}$ is a curve as the image by $(\nabla f, f)$ of the polar curve $X := \nabla f^{-1}(\mathbb{C}w)$ (∇f is surjective and we are in a neighbourhood of the origin). Fix (w, t) and denote by Γ an irreducible component of $(\nabla f, f)(X)$ for which $(w, t) \in C_0(\Gamma)$. Then there exists $v \in \mathbb{C}^{m+1} \setminus \{0\}$ such that $C_0(\Gamma) = \mathbb{C}v$. Now we need the following lemma.

Lemma 2.2. *Let $h: W \rightarrow \mathbb{C}^n$ be a holomorphic function defined in a neighbourhood $W \subset \mathbb{C}$ of zero, satisfying $h(0) = 0$ and such that $h(W)$ defines an irreducible curve germ at zero. Then there is a sequence $W \ni t_\nu \rightarrow 0$ such that $h(t_\nu)/\|h(t_\nu)\|$ converge to a vector v for which $C_0(h(W)) = \mathbb{C}v$.*

Proof of the Lemma. Take a sequence $W \setminus h^{-1}(0) \ni t_\nu \rightarrow 0$. Since $\{h(t_\nu)/\|h(t_\nu)\|\} \subset \mathbb{S}^{2n-1}$, extracting, if necessary, a subsequence, we may assume that these terms converge to some $v \in \mathbb{S}^{2n-1}$. But $h(t_\nu) \rightarrow 0$ and $\lambda_\nu = 1/\|h(t_\nu)\| > 0$. Hence $v \in C_0(h(W))$, whence the cone contains the line $\mathbb{C}v$. The germ at zero of $h(W)$ being irreducible, the tangent cone coincides with the line found. $\square$

Let $\Gamma' \subset X$ be an irreducible component such that $(\nabla f, f)(\Gamma') = \Gamma$. Consider a Puiseux parametrization $\gamma: W \rightarrow \Gamma'$. Then $h := (\nabla f, f) \circ \gamma$ satisfies the assumptions of the preceding lemma and so we have a sequence $t_\nu \rightarrow 0$ such that $v = \lim h(t_\nu)/\|h(t_\nu)\|$. Write $v = (v', v'') \in \mathbb{C}^m \times \mathbb{C}$; we will show that $v'' = 0$. We look at $f(\gamma(t_\nu))$; on the one hand,

$$|f(\gamma(t_\nu))| \leq \text{const.} |t_\nu|^{\text{ord}_0(f \circ \gamma)}, \quad \nu \gg 1,$$

whereas on the other,

$$\|h(t_\nu)\| \geq \|\nabla f(\gamma(t_\nu))\| \geq \text{const.} |t_\nu|^{\text{ord}_0(\nabla f \circ \gamma)}, \quad \nu \gg 1,$$

where obviously⁴ $\text{ord}_0(\nabla f \circ \gamma) = \min_j \text{ord}_0(\frac{\partial f_j}{\partial z} \circ \gamma)$. For ν large enough,

$$\frac{|f(\gamma(t_\nu))|}{\|h(t_\nu)\|} \leq \text{const.} |t_\nu|^{\text{ord}_0(f \circ \gamma) - \text{ord}_0(\nabla f \circ \gamma)}$$

and it remains to observe that the exponent is positive.

To see this we argue classically as follows: if $g: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ is an analytic germ, then $\text{ord}_0 g = \text{ord}_0 g' + 1$, whence $\text{ord}_0(f \circ \gamma) =$

⁴From the expansion into a power series it is easy to see that for a non-constant analytic germ $g: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ there is $c_1|t|^q \leq |g(t)| \leq c_2|t|^q$ in a neighbourhood of zero with $q = \text{ord}_0 g$, $c_1, c_2 > 0$. Hence, if $g = (g_1, \dots, g_n)$ on $(\mathbb{C}, 0)$ the same kind of inequality holds true with $q = \min_j \text{ord}_0 g_j$.

$\text{ord}_0(f \circ \gamma)' + 1$. But (since $\text{ord}_0 \sum_1^k g_j \geq \min_j \text{ord}_0 g_j$)

$$\begin{aligned} \text{ord}_0(f \circ \gamma)' &= \text{ord}_0 \langle \nabla f \circ \gamma, \gamma' \rangle \geq \\ &\geq \min_j \text{ord}_0 \left(\frac{\partial f}{\partial z_j} \circ \gamma \right) \gamma'_j = \\ &= \min_j \left\{ \text{ord}_0 \left(\frac{\partial f}{\partial z_j} \circ \gamma \right) + \text{ord}_0 \gamma'_j \right\}. \end{aligned}$$

Since $\text{ord}_0 \gamma'_j \geq 0$ for all j , the last expression is greater or equal to $\text{ord}_0(\nabla f \circ \gamma)$.

Summing up, we have proved the following Proposition (cf. [T] Corollaire 2 p. 270 and [P1], [PW]):

Proposition 2.3. *If $f \in \mathcal{O}_m$ is a holomorphic germ such that ∇f has an isolated zero at the origin, then the coefficients of the characteristic polynomial $P(w, t) = t^\mu + a_1(w)t^{\mu-1} + \dots + a_\mu(w) \in \mathcal{O}_m[t]$, of f with respect to ∇f , where μ is the Milnor number of f , all satisfy $\text{ord}_0 a_j \geq j + 1$ and we have in a neighbourhood of zero the gradient inequality*

$$|f(z)| \leq \text{const.} \cdot \|\nabla f(z)\|^{\max_{j=1}^\mu \frac{j}{\text{ord}_0 a_j}}.$$

3. GRADIENT INEQUALITY WITH PARAMETER

Let $f(x, t) \in \mathcal{O}_{m+k}$ be a holomorphic germ. Write $f_t(x) := f(x, t)$ and put $g_t(x) := \nabla f_t(x)$ for the complex gradient. The resulting germ $g(x, t) := g_t(x)$ is, of course, holomorphic.

We assume that for all t small enough, $g_t^{-1}(0) = \{0\}$ is isolated at zero (note that this implies that $f(0, t) \equiv 0$ e.g. by the gradient inequality) and that the Milnor numbers $m_0(g_t) \equiv \mu$ are constant.

The theorem we are aiming at is the following:

Theorem 3.1. *If $f(x, t) \in \mathcal{O}_{m+k}$ is such that for all t small enough, $f_t(x)$ has an isolated singularity at zero with constant Milnor number μ , then for all (x, t) in a neighbourhood of zero*

$$|f_t(x)|^{\frac{\mu}{\mu+1}} \leq C \|\nabla f_t(x)\|$$

where $C > 0$. Note that the exponent belongs to $[1/2, 1)$ and f itself satisfies the gradient inequality (with respect to (x, t)) with this exponent.

The proof will be given in several steps.

Step 1

Consider the germ $G(x, t) := (g(x, t), t)$. Since $G^{-1}(0) = g_0^{-1}(0) \times \{0\}$, $G: (\mathbb{C}^{m+k}, 0) \rightarrow (\mathbb{C}^{m+k}, 0)$ is the germ of a proper mapping.

Lemma 3.2. *In the setting introduced, the multiplicity at zero of the branched covering G is μ .*

Proof. We have $G^{-1}(0) = \{0\}$. If (y, t) is a point close enough to zero but not belonging to the critical locus, then $m_0(G) = \#G^{-1}(y, t)$. On the other hand, $G^{-1}(y, t) = g_t^{-1}(y) \times \{t\}$, whence

$$m_0(G) \leq \#g_t^{-1}(y) \leq \mu.$$

However, if y does not lie in the critical locus of g_0 , then

$$\mu = \#g_0^{-1}(y) = \#G^{-1}(y, 0) \leq m_0(G).$$

□

Step 2

Let now $P(y, t, s) \in \mathcal{O}_{m+k}[s]$ be the characteristic polynomial of f with respect to G . Therefore, we have $P(t, g(x, t), f(x, t)) \equiv 0$ and so by the Viète formulæ:

$$|f(x, t)| \leq 2 \max_{j=1}^{\mu} |a_j(g(x, t), t)|^{1/j}$$

where we write $P(y, t, s) = s^{\mu} + a_1(y, t)s^{\mu-1} + \dots + a_{\mu}(y, t)$.

Lemma 3.3. $P(\cdot, t, \cdot)$ is the characteristic polynomial of f_t with respect to g_t .

Proof. For a generic point y near zero there is $\#g_t^{-1}(y) = \mu$. Let $x^{(1)}, \dots, x^{(\mu)}$ be the points in this fibre. Then since the fibre $G^{-1}(y, t)$ consists precisely of the points $(x^{(j)}, t)$, it is maximal and we get

$$P(y, t, s) = \prod_{j=1}^{\mu} (s - f(x^{(j)}, t)) = \prod_{j=1}^{\mu} (s - f_t(x^{(j)}))$$

which gives the assertion. □

In particular we have $a_j(0, t) = 0$.

Step 3

Fix j and consider the expansion of a_j into a Hartogs series in a polydisc $P(r)$ centred at zero (the same for all $j = 1, \dots, \mu$)

$$a_j(y, t) = \sum_{\nu=\nu_j(t)}^{+\infty} \sum_{\substack{\alpha \in \mathbb{Z}_+^m \\ |\alpha|=\nu}} \left(\frac{1}{\alpha!} D^{(\alpha,0)} a_j(0, t) \right) y^{\alpha}$$

where $\nu_j(t) := \text{ord}_0 a_j(\cdot, t) \geq j + 1$ by Proposition 2.3. The Cauchy inequalities yield

$$\left| \frac{1}{\alpha!} D^{(\alpha,0)} a_j(0, t) \right| \leq \frac{1}{r^{|\alpha|}} M_j$$

with $M_j := \sup\{|a_j(y, t)| : (y, t) \in P(r)\} > 0$. Let $M = \max_{j=1}^{\mu} M_j$. Writing $j+1$ instead of $\nu_j(t)$ we have in $P(r)$,

$$\begin{aligned} |a_j(y, t)| &\leq \sum_{\nu=j+1}^{+\infty} \frac{M}{r^{\nu}} \|y\|^{\nu} \leq \\ &\leq \|y\|^{j+1} \frac{M}{r^{j+1}} \sum_{\nu=0}^{+\infty} \left(\frac{\|y\|}{r} \right)^{\nu}. \end{aligned}$$

This leads to the conclusion that for some constant $C > 0$ we have for all (y, t) in a neighbourhood V of zero and all indices j ,

$$|a_j(y, t)| \leq C \|y\|^{j+1}.$$

Step 4

Note that $g(0, t) = 0$ and we obtain the inequalities

$$|a_j(g(x, t), t)| \leq c_j(t) \|g(x, t)\|^{j+1}.$$

for all j and (x, t) in a neighbourhood U of zero such that $G(U) \subset V$.

This gives the gradient inequality with parameter and

$$\max_{j=1}^{\mu} \frac{j}{j+1} = \frac{\mu}{\mu+1}$$

as exponent. The proof is complete.

4. ACKNOWLEDGEMENTS

This research was partially supported by Polish Ministry of Science and Higher Education grant ...

REFERENCES

- [Ch] E. M. Chirka, *Complex Analytic Sets*, Kluwer Acad. Publ. 1989;
- [Ł] S. Łojasiewicz, *Sur les trajectoires du gradient d'une fonction analytique*, Sem. di Geom. 1982-1983, Univ. di Bologna (1983), pp. 115-117;
- [P1] A. Płoski, *Multiplicity and the Łojasiewicz exponent*, Banach Centre Publ. 20 - Singularities, Warsaw 1985 (1988), pp. 353-354;
- [P2] A. Płoski, *Semicontinuity of the Łojasiewicz exponent*, Univ. Iagel. Acta Math. No. 48 (2010), 103-110;
- [PW] A. Płoski, T. Winiarski, *On a theorem of Platte*, Bull. Polish Acad. Sci. 39 nr 1-2, (1991), 57-61;
- [T] B. Teissier, *Variétés polaires I: Invariants polaires des singularités d'hypersurfaces*, Invent. Math. 40 (1977), 267-292.

JAGIELLONIAN UNIVERSITY, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, INSTITUTE OF MATHEMATICS, UL. ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

E-mail address: maciej.denkowski@uj.edu.pl